

AESB1320-17 - Solutions

11-04-18, 9:00

TN-TZ 4.25

The exam consists of 10 Conceptual Questions (CQs), each valid for 1 point, and 10 Exercises (EXs), each valid for 4 points. The maximum score is 50. The pass score is 29.

PART I: all students.

CQ1

A pilot drops a package from a plane flying horizontally at a constant speed. Neglecting air resistance, when the package hits the ground the horizontal location of the plane will

- A. be in front of the package.
- B. be behind the package.
- C. depend of the speed of the plane when the package was released.
- D. be over the package.

D: there is no force acting in the horizontal direction (air resistance is neglected), so plane and package will keep moving forward with the same velocity.

CQ2

Three people are pushing a car up a hill at constant velocity. The net force on the car is

- A. zero.
- B. up the hill and greater than the weight of the car.
- C. down the hill and greater than the weight of the car.
- D. down the hill and equal to the weight of the car.
- E. up the hill and equal to the weight of the car.

A: If the car moves with constant velocity, then the net acceleration has to be zero.

CQ3

Which of the following statements is true?

- a. The total mechanical energy of a system is constant only if non-conservative forces act.
- b. The total mechanical energy of a system, at any one instant, is either all kinetic or all potential energy.
- c. The total mechanical energy of a system is constant only if conservative forces act.
- d. The total mechanical energy of a system is equally divided between kinetic and potential energy.

c: According to the definition of conservative force.

CQ4

If the mass of the earth and all objects on it were suddenly doubled, but the size remained the same, the acceleration due to gravity at the surface would become

- a. 2 times what it now is.
- b. the same as it now is.
- c. 4 times what it now is.
- d. 1/2 of what it now is.
- e. 1/4 of what it now is.

a: $g=GM/R^2$, so doubling M doubles g .

CQ5

There must be equal amounts of mass on both side of the center of mass of an object.

- a. True
- b. False

b: The position of the center of mass depends on both the amount of mass and its spatial distribution (i.e., you can have more mass at one side, if it's closer).

CQ6

If two forces of equal magnitude act on an object that is hinged at a pivot, the force acting farther from the pivot must produce the greater torque about the pivot.

- A. false
- B. true
- C. unable to decide without knowing the shape of the object

A: It also depends on the angle between the force and the line between the pivot and the application point of the force itself.

CQ7

When you ride a bicycle, in what direction is the angular velocity of the wheels?

- A. to your right
- B. to your left
- C. backwards
- D. up
- E. forwards

B: Right-hand rule.

EX1

A hockey puck slides off the edge of a table with an initial velocity of 27.7 m/s and experiences no air resistance. The height of the tabletop above the ground is 2.00 m.

What is the angle below the horizontal of the velocity of the puck just before it hits the ground?

12.7°

Data:

$$v_{0x} = v_0 = 27.7 \text{ m/s}; y_0 = h = 2.0 \text{ m}; \theta = ?$$

Solution:

Note: the question is about instantaneous velocity before hitting the ground, not average quantities. Horizontal velocity remains unchanged (v_{0x}), vertical velocity (v_y) is the velocity a body in free fall has after falling vertically by 2 meters.

Since you are not asked for the travel time, you can use:

$$v_y^2 = v_0^2 - 2g(y - y_0) = 2gh \Rightarrow v_y = \pm \sqrt{2gh} = -6.26 \text{ m/s} \text{ (positive sign also ok, if y axis is downward)}$$

Alternatively, you can first solve for time using: $y = y_0 + v_{0y}t - 1/2gt^2 \Rightarrow t = \sqrt{2h/g} = 0.64 \text{ s}$

And then use: $v_y = v_{0y} - gt \Rightarrow v_y = -gt = -6.27 \text{ m/s}$

The angle is just given by: $\theta = \arctan(v_y/v_x) = 12.7^\circ$ (or -12.7 , if you measure angles counter-clockwise).

EX2

You push downward on a box at an angle 25° below the horizontal with a force of 750 N. The box is on a flat horizontal surface for which the coefficient of static friction with the box is 0.64.

What is the mass of the heaviest box you will be able to move?

76 kg

Data:

$$\theta = 25^\circ; F = 750 \text{ N}; \mu = 0.64; m_{\max} = ?$$

Solution:

Friction will equal the total normal (vertical) force, which is equal to the weight of the box, plus the vertical component of the force pushing down the box.

$$\text{Vertical component of the pushing force: } F_y = F \sin \theta = 315 \text{ N}$$

$$\text{Horizontal component of the pushing force: } F_x = F \cos \theta = 679.7 \text{ N}$$

$$\text{Force of friction: } F_f = \mu n = \mu (F_y + mg)$$

So, the maximum mass that can be moved is the one for which the force of friction equals the horizontal component of the pushing force:

$$F_f = F_x \Rightarrow \mu (F \sin \theta + mg) = F \cos \theta \Rightarrow m_{\max} = F (\cos \theta - \mu \sin \theta) / (\mu g) = 76 \text{ kg}$$

EX3

A very small 100-g object is attached to one end of a massless 10-cm rod that is pivoted without friction about the opposite end. The rod is held vertical, with the object at the top, and released, allowing the rod to swing. What is the speed of the object at the instant that the rod is horizontal?

1.4 m/s

Data:

$$m = 100 \text{ g}; h = 0.1 \text{ m}; v(\text{horiz}) = ?$$

Solution:

$$\text{At the top, all energy is potential: } E_0 = mgh = 0.098 \text{ J}$$

$$\text{When horizontal, all energy is kinetic: } E_1 = 1/2 mv^2$$

$$\text{Since the only force is gravity, energy is conserved: } E_0 = E_1 \Rightarrow v = \sqrt{2gh} = 1.4 \text{ m/s}$$

$$\text{Also possible to start from rotational kinetic energy: } E_1 = 1/2 I\omega^2 = 1/2 mh^2 (v/h)^2 = 1/2 mv^2$$

EX4

Three identical very small 50-kg masses are held at the corners of an equilateral triangle, 0.30 m on each side. If one of the masses is released, what is its initial acceleration if the only forces acting on it are the gravitational forces due to the other two masses? ($G = 6.67 \times 10^{-11} \text{ N m}^2/\text{kg}^2$)

$6.4 \times 10^{-8} \text{ m/s}^2$

Data:

$$m = 50 \text{ kg}; d = 0.3 \text{ m}; a_i = ?$$

Solution:

$$\text{Equilateral triangle: } \theta = 60^\circ$$

$$\text{In horizontal direction: } F_{1x} + F_{2x} = 0$$

$$\text{In vertical direction: } F_{1y} = F_{2y} = Gmm/d^2 \cdot \cos(\theta/2) \Rightarrow a = (F_{1y} + F_{2y})/m = 2Gm/d^2 \cdot \cos 30^\circ = 6.4 \times 10^{-8} \text{ m/s}^2$$

EX5

A billiard ball traveling at 3.00 m/s collides perfectly elastically with an identical billiard ball initially at rest on the level table. The initially moving billiard ball deflects 30.0° from its original direction. What is the speed of the initially stationary billiard ball after the collision?

1.55 m/s

Total linear momentum (of both balls together) is conserved: it remains constant in the direction of initial motion and zero in the perpendicular direction.

$$p_x = mv_0 = mv_{1x} + mv_{2x} \Rightarrow v_{2x} = v_0 - v_{1x} = v_0(1 - \cos 30^\circ) = 0.4 \text{ m/s}$$

$$p_y = mv_{1y} + mv_{2y} = 0 \Rightarrow v_{2y} = -v_{1y} = -v_0 \sin 30^\circ = -1.5 \text{ m/s}$$

$$v_2 = \sqrt{v_{2x}^2 + v_{2y}^2} = \sqrt{2.41} = 1.55 \text{ m/s}$$

EX6

An extremely light rod 1.00 m long has a 2.00-kg mass attached to one end and a 3.00-kg mass attached to the other. The system rotates at a constant angular speed about a fixed axis perpendicular to the rod that passes through the rod 30.0 cm from the end with the 3.00-kg mass attached. The kinetic energy of the system is measured to be 100.0 J.

What is the moment of inertia of this system about the fixed axis?

$$1.25 \text{ kg m}^2$$

Since the shape of the masses is not given, the fact that the masses are small compared to the length of the rod should have been mentioned (point masses). If this motivation led to the answer "impossible to determine", it will be accepted.

Solution:

The moment of inertia only depends on the position of the two masses about the rotation axis:

$$I = m_1 r_1^2 + m_2 r_2^2$$

EX7

A 5.0-m radius playground merry-go-round with a moment of inertia of 2000 kg m² is rotating freely with an angular speed of 1.0 rad/s. Two people, each having a mass of 60 kg, are standing right outside the edge of the merry-go-round and step on it with negligible speed. What is the angular speed of the merry-go-round right after the two people have stepped on?


$$0.4 \text{ rad/s}$$

Total angular momentum (merry-go-round) is conserved.

$$I_{\text{people}} = 2mR^2 = 3000 \text{ kg m}^2$$

$$L = \omega_0 I_{\text{merry-go-round}} = \omega_1 (I_{\text{merry-go-round}} + I_{\text{people}}) \Rightarrow \omega_1 = \omega_0 I_{\text{merry-go-round}} / (I_{\text{merry-go-round}} + I_{\text{people}}) = 0.4 \text{ rad/s}$$

EX8



A 10.0-kg uniform ladder that is 2.50 m long is placed against a smooth vertical wall and reaches to a height of 2.10 m, as shown in the figure. The base of the ladder rests on a rough horizontal floor whose coefficient of static friction with the ladder is 0.800. An 80.0-kg bucket of concrete is suspended from

the top rung of the ladder, right next to the wall, as shown in the figure. What is the magnitude of the friction force that the floor exerts on the ladder?

539.5 N

Data:

$m_L=10$ kg; $L_L=2.5$ m; $h=2.1$ m; $\mu=0.8$; $m_B=80$ kg; $F_f=?$

Additional: $d=\sqrt{L_L^2-h^2}=1.36$ m

Solution 1 (pivot point at the base of the ladder, point O):

Horizontal forces: $-N_w+F_f=0 \Rightarrow F_f=N_w$

with N_w being the normal force by the wall (horizontal towards the left)

Torque about O: $-m_Bgd+N_w h-m_Lgd/2=0 \Rightarrow N_w=gd(m_B+m_L/2)/h=539.5$ N

Solution 2 (pivot point at the top of the ladder, point P):

Vertical forces: $N_F-m_Lg-m_Bg=0 \Rightarrow N_F=g(m_L+m_B)$

with N_F being the normal force by the floor (vertical upwards)

Torque about P: $F_f h-N_F d+m_Lgd/2=0 \Rightarrow F_f=gd(m_L+m_B-m_L/2)/h=539.5$ N

PART II: *only first-year students.*

CQ8-a

CQ9-a

CQ10-a

EX9-a

EX10-a

PART III: *only students who take the resit of Mechanics 1.*

CQ8-b

If two vectors point in opposite directions, their cross product must be zero.

- True
- False

A: The magnitude of the cross product depends on $\sin(\theta)$, and $\sin 180^\circ=0$.

CQ9-b

A box of mass m is pulled with a constant acceleration ' a ' along a horizontal frictionless floor by a wire that makes an angle of 15° above the horizontal. If T is the tension in this wire, then

- $T < ma$.
- $T = ma$.

c. $T > ma$.

c: You need ma to accelerate the box horizontally, plus a vertical component due to the fact that the wire is not horizontal (note that the vertical component does not affect the acceleration).


CQ10-b

If a force always acts perpendicular to an object's direction of motion, that force cannot change the object's kinetic energy.

- A. True
- B. False

A: The force will not change the magnitude of the velocity, so the kinetic energy remains constant.

EX9-b



Two blocks are connected by a string that goes over an ideal pulley as shown in the figure. Block A has a mass of 3.00 kg and can slide over a rough plane inclined 30.0° to the horizontal. The coefficient of kinetic friction between block A and the plane is 0.40. Block B has a mass of 2.77 kg. What is the acceleration of the blocks?

0.374

The two blocks have almost the same mass, but A is on a small slope, so the force of gravity on block B is larger and block B will fall (and block A move up). On block A you have the component of gravity pointing down the slope, friction also pointing down the slope (because A moves up), and the tension T pointing up. On block B you only have gravity and tension. The acceleration of the two blocks is the same, since the pulley has no mass.

Block A: $m_A a = -m_A g \sin 30^\circ - m_A g \cos 30^\circ \mu + T$ [0.5 points if correct]

Block B: $m_B a = m_B g - T$ [0.5 points if correct]

Solving for acceleration (a) you obtain:

$$a = g [-m_A (\sin 30^\circ - \cos 30^\circ \mu) + m_B] / (m_A + m_B) = 9.81 * (0.22 / 5.77) = 0.374 \text{ m/s}^2$$

EX10-b

A ball is thrown upward at an angle with a speed and direction such that it reaches a maximum height of 16.5 m above the point it was released, with no appreciable air resistance. At its maximum height it has a speed of 15.0 m/s. With what speed was the ball released?

23.4 m/s

Only force is gravity (=conservative), which acts in vertical direction.

Conservation of mechanical energy in y direction: initial kinetic is equal to final potential:

$$1/2 m (v_{0y})^2 = mgd, \text{ with } d = 16.5 \text{ m}$$

In x direction v is constant, so it is equal to the value at maximum height: $v_{0x} = 15 \text{ m/s}$.

Since energy is conserved, the initial velocity is the squared sum of the two components.